

Generalization of an example of Hartshorne concerning local cohomology

M. Hellus and J. Stückrad

February 2, 2008

Abstract

We prove the following generalization of an example of Hartshorne: Let k be a field, $n \geq 4$, $R = k[[X_1, \dots, X_n]]$, $I = (X_1, \dots, X_{n-2})R$ and $p \in R$ a prime element such that $p \in (X_{n-1}, X_n)R$. Then $H_I^{n-2}(R/pR)$ is not artinian.

1 Introduction

It is an interesting question to determine if a given local cohomology module $H_I^i(M)$ is artinian, where I is an ideal of a noetherian local ring (R, m) and M is a finite R -module; in fact, this is one of Huneke's problems on local cohomology (see [8, third problem]).

In general a given R -module A is artinian if and only if both

$$\text{Supp}_R(A) \subseteq \{m\} \quad (A_1)$$

and

$$\text{soc}(A) := \text{Hom}_R(R/m, A) \text{ is finitely generated} \quad (A_2)$$

hold. Hartshorne (see [2, section 3]) showed that $H_I^2(R)$ is not artinian, where k is a field, $R = k[[u, v, x, y]]/(ux + vy)$ and $I \subseteq R$ is the ideal generated by the classes of u and v in R (in fact Hartshorne showed something slightly different, but it is not difficult to modify his result to the statement mentioned above). More precisely, $H_I^2(R)$ fulfills (A_1) , but not (A_2) .

On the other hand, recently, modules of the form $D(H_I^i(M))$ have been studied for general R , I , i and M (see [3, 4, 5, 6, 7], $D(_)$ is the Matlis dual functor, see [9] for details on Matlis duality). It turns out that the study of these objects leads to an elegant proof of Hartshorne's example; in fact we show that $D(H_I^2(R))$ is not noetherian, where $H_I^2(R)$ is the local cohomology module of Hartshorne's example. Hence $H_I^2(R)$ cannot be artinian. Furthermore, our method can be used to show the following more general result:

Theorem 1.1. *Let k be a field, $R = k[[X_1, \dots, X_n]]$ a power series algebra over k ($n \geq 3$), $I = (X_1, \dots, X_{n-2})R$ and p a prime element of R such that $p \in (X_{n-1}, X_n)R$. Then $H_I^{n-2}(R/pR)$ fulfills (A_1) , but not (A_2) ; in particular, it is not artinian.*

It should be remarked that Marley and Vassilev (see [10, theorem 2.3]) have generalized Hartshorne's example in a different direction. Due to different hypothesis, their and our generalization can be compared only in a special case, see remark 2.6 for details.

The authors thank Gennady Lyubeznik for drawing their attention to Hartshorne's example.

2 Results

Let k be a field, $n \geq 3$, $R = k[[X_1, \dots, X_n]]$ a formal power series algebra over k , I the ideal $(X_1, \dots, X_{n-2})R$ of R and define

$$D := D(H_I^{n-2}(R)) \quad ,$$

where $D(-)$ is the Matlis dual functor, i. e. $D(M) := \text{Hom}_R(M, E_R(R/m))$ for any R -module M , $E_R(R/m)$ being a fixed R -injective hull of R/m . Our method is to study the module D , here are some properties of D :

Properties 2.1. (i) *Every associated prime $p \in \text{Ass}_R(D)$ has $\text{height}(p) \leq 2$.*
(ii) *For every prime ideal p of R with $\text{height}(p) = 2$ one has:*

$$p \in \text{Ass}_R(D) \iff I + p \text{ is } m\text{-primary}.$$

Proof. (i) Let $p \in \text{Ass}_R(D)$. We conclude

$$\begin{aligned} 0 &\neq \text{Hom}_R(R/p, D) \\ &= D(H_I^{n-2}(R)/pH_I^{n-2}(R)) \\ &= D(H_I^{n-2}(R/p)) \quad . \end{aligned}$$

Here the first equality follows formally from the exactness of $D(-)$ and the second from the right exactness of the functor $H_I^{n-2}(-) = H_{(X_1, \dots, X_{n-2})R}^{n-2}(-)$.

It is well-known that $H_I^{n-2}(R/p) \neq 0$ implies $\dim(R/p) \geq n - 2$.

(ii) Let p be a prime ideal of R of height two. Because of (i), we have

$$p \in \text{Ass}_R(D) \iff \text{Hom}_R(R/p, D) \neq 0 \quad .$$

In the proof of (i) it was shown that the latter module equals

$$D(H_I^{n-2}(R/p))$$

and, by the well-known Hartshorne-Lichtenbaum vanishing theorem, is non-zero if and only if the ideal $I + p$ is m -primary.

Example 2.2. *In the above situation, take $n = 4$ and, for every $\lambda \in k$, define*

$$p_\lambda := (X_3 + \lambda X_1, X_4 + \lambda X_2)R \quad .$$

Clearly, every p_λ is a height two prime ideal of R and, by property (ii) above, is associated to $D = D(H_I^2(R))$. On the other hand, for every $\lambda \in k$, one has

$$p := X_1 X_4 + X_2 X_3 \in p_\lambda$$

(because of $p = X_1(X_4 - \lambda X_2) + X_2(X_3 + \lambda X_1)$). Therefore, at least if k is infinite, D has infinitely many associated primes containing p . This implies that

$$\text{Hom}_R(R/pR, D)$$

cannot be finitely generated. But, as we have seen in the proof of property (i) above, $\text{Hom}_R(R/pR, D)$ is the Matlis dual of

$$H_I^2(R/pR)$$

and so $H_I^2(R/pR)$ cannot be artinian.

Remark 2.3. *This is essentially Hartshorne's example ([2, section 3]), the main difference is that Hartshorne works over the ring $k[X_3, X_4][[X_1, X_2]]$, while we work over the ring $k[[X_1, X_2, X_3, X_4]]$; but the two versions are essentially the same, because the module*

$$H_{(X_1, X_2)}^2(k[X_3, X_4][[X_1, X_2]]/(X_1 X_4 + X_2 X_3))$$

is naturally a module over $k[[X_1, X_2, X_3, X_4]]$, because its support is $\{(X_1, X_2, X_3, X_4)\}$. This is true, because for every prime ideal p of $k[X_3, X_4][[X_1, X_2]]$ different from (X_1, X_2, X_3, X_4) and containing $X_1 X_4 + X_2 X_3$ the ring $(k[X_3, X_4][[X_1, X_2]]/(X_1 X_4 + X_2 X_3))_p$ is regular, and so Hartshorne-Lichtenbaum vanishing shows that

$$H_{(X_1, X_2)}^2(k[X_3, X_4][[X_1, X_2]]/(X_1 X_4 + X_2 X_3))_p = 0 \quad .$$

A similar technique like in example 2.2 works to show that $H_I^{n-2}(R/pR)$ is not artinian for $R = k[[X_1, \dots, X_n]]$, $n \geq 4$, $p \in (X_{n-1}, X_n)R$ a prime element and k an arbitrary field (may be finite):

Theorem 2.4. *Let k be a field, $n \geq 4$, $R = k[[X_1, \dots, X_n]]$, $I = (X_1, \dots, X_{n-2})R$ and $p \in R$ a prime element such that $p \in (X_{n-1}, X_n)R$. Set $D := D(H_I^{n-2}(R))$.*

(i) If $p \in R$ is a prime element such that $p \in (X_{n-1}, X_n)R \cap I$ holds, the set

$$\{p \in \text{Ass}_R(D) | p \in p\}$$

is infinite.

(ii) If $p \in R$ is a prime element such that $p \in (X_{n-1}, X_n)R$, $H_I^{n-2}(R/pR)$ is not artinian.

Proof. (i) It is easy to see that there exist $f, g \in I, f \notin (X_{n-1}, X_n)R$ and $l \geq 1$ such that

$$p = X_n^l f + X_{n-1} g$$

holds (note that f is not zero because p is prime). Let $m \in \mathbb{N}^+$ be arbitrary. We have

$$p = (X_n^l + X_1^m g) f + (X_{n-1} - X_1^m f) g$$

and so $p \in I_m := (X_n^l + X_1^m g, X_{n-1} - X_1^m f)R$. The elements $X_1, \dots, X_{n-2}, X_n^l + X_1^m g, X_{n-1} - X_1^m f$ form a system of parameters of R and so, by properties (i) and (ii) from above, there exists a $p_m \in \text{Ass}_R(D(H_I^{n-2}(R)))$ containing I_m . p_m necessarily has height two. For $m, m' \in \mathbb{N}^+, m \neq m'$

$$\sqrt{I_m + I_{m'}} = (X_1, X_n, X_{n-1})R \cap \sqrt{(X_{n-1}, X_n, f, g)R}$$

holds; in particular, all primes containing $I_m + I_{m'}$ have height at least three. The statement follows now from property (i).

(ii) If $p \notin I$, it is easy to see that

$$\text{Supp}_R(H_I^{n-2}(R/pR)) = \mathcal{V}(I + pR) ,$$

the set of prime ideals of R containing $I + pR$, and so $H_I^{n-2}(R/pR)$ does not satisfy (A_1) . We assume $p \in I$: If $H_I^{n-2}(R/pR)$ was artinian, its dual $D(H_I^{n-2}(R/pR))$ would be finitely generated; but we have seen before that, because of the exactness of D and the right-exactness of $H_I^{n-2}(-)$, $D(H_I^{n-2}(R/pR)) = \text{Hom}_R(R/pR, D)$, and from (i) we know that the latter module is not finitely generated.

Remark 2.5. *Marley and Vassilev have shown*

Theorem ([10, theorem 2.3])

Let (T, m) be a noetherian local ring of dimension at least two. Let $R = T[x_1, \dots, x_n]$ be a polynomial ring in n variables over T , $I = (x_1, \dots, x_n)$, and $f \in R$ a homogenous polynomial whose coefficients form a system of parameters for T . Then the $^\text{socle}$ of $H_I^n(R/fR)$ is infinite dimensional.*

In their paper [10], Marley and Vassilev say (in section 1) that Hartshorne's example is obtained by letting $T = k[[u, v]]$, $n = 2$ and $f = ux + vy$; there is a slight difference between the two situations that comes from the fact that Hartshorne works over a ring of the form $k[x, y][[u, v]]$ while Marley and Vassilev work over a ring of the form $k[[u, v]][x, y]$. The two rings are not the same. But, as

$$\mathrm{Supp}_R(H_{(u,v)}^2(R/(uy + vx))) = \{(x, y, u, v)\}$$

(both for $R = k[x, y][[u, v]]$ and for $R = k[[u, v]][x, y]$), the local cohomology module in question is (in both cases) naturally a module over $k[[x, y, u, v]]$ and, therefore, both versions are equivalent, i. e. the result of Marley and Vassilev is a generalization of Hartshorne's example.

Remark 2.6. [10, theorem 2.3] and our theorem 2.4 are both generalizations of Hartshorne's example, but, due to different hypotheses, they can only be compared in the following special case: k a field, $n \geq 4$,

$$R_0 = k[[X_{n-1}, X_n]][X_1, \dots, X_{n-2}] ,$$

$$R = k[[X_1, \dots, X_n]] ,$$

$I = (X_1, \dots, X_{n-2})R$, $p \in R_0$ a homogenous element such that p is prime as an element of R . Then [10, theorem 2.3] says (implicitly) that

$$H_I^{n-2}(R/pR)$$

is not artinian, if the coefficients of $p \in R_0$ in $k[[X_{n-1}, X_n]]$ form a system of parameters in $k[[X_{n-1}, X_n]]$, while theorem 2.4 says that the same module is not artinian if none of these coefficients of p is a unit in $k[[X_{n-1}, X_n]]$.

References

- [1] Grothendieck, A. Local Cohomology, *Lecture Notes in Mathematics*, Springer Verlag, (1967).
- [2] Hartshorne, R. Affine Duality and Cofiniteness, *Invent. Math.* **9**, (1970) 145–164.
- [3] Hellus, M. On the associated primes of Matlis duals of top local cohomology modules, *Communications in Algebra* **33**, (2005), no. 11, 3997–4009.
- [4] Hellus, M. Matlis duals of top local cohomology modules and the arithmetic rank of an ideal, to appear in *Communications in Algebra*.
- [5] Hellus, M. Attached primes and Matlis duals of local cohomology modules, to appear in *Archiv der Mathematik*

- [6] Hellus, M. Finiteness properties of duals of local cohomology modules, to appear in *Communications in Algebra*.
- [7] Hellus, M. and Stückrad, J. Matlis duals of top Local Cohomology Modules, submitted to *Proceedings of the American Mathematical Society*.
- [8] Huneke, C. Problems on Local Cohomology, *Res. Notes Math.* **2**, (1992) 93–108.
- [9] Matlis, E. Injective modules over Noetherian rings, *Pacific J. Math.* **8**, (1958) 511–528.
- [10] Marley, M. and Vassilev, J.C. Local cohomology modules with infinite dimensional socles, *Proc. Amer. Math. Soc.* Vol. **132**, No **12**, (2004) 3485–3490.

UNIVERSITÄT LEIPZIG, FAKULTÄT FÜR MATHEMATIK UND INFORMATIK,
 MATHEMATISCHES INSTITUT, AUGUSTUSPLATZ 10/11, D-04109 LEIPZIG
E-mail:

hellus@math.uni-leipzig.de
 stueckrad@math.uni-leipzig.de